

ASN Neuro, 2009, Aug 14;1(3)

Cuprizone demyelination of the corpus callosum in mice correlates with altered social interaction and impaired bilateral sensorimotor co-ordination

Norah Hibbits*, Ravinder Pannu[†], T John Wu[‡] and Regina C Armstrong*[†]

*Program in Neuroscience, Uniformed Services University of the Health Sciences, Bethesda, MD 20814, U.S.A.

[†]Department of Anatomy, Physiology and Genetics, Uniformed Services University of the Health Sciences, Bethesda, MD 20814, U.S.A.

[‡]Department of Obstetrics and Gynecology, Uniformed Services University of the Health Sciences, Bethesda, MD 20814, U.S.A.

Experimental Neurology 2007 May;205(1): 207–213

A highly sensitive automated complex running wheel test to detect latent motor deficits in the mouse MPTP model of Parkinson's disease

David Liebetanz^a, Paul Christian Baier^a, Walter Paulus^a, Katrin Meuer^{b,c},
Mathias Bähr^{b,c}, Jochen H. Weishaupt^{b,c}

^a Department of Clinical Neurophysiology, Georg-August-University, Robert-Koch Strasse 40, 37099 Göttingen, Germany

^b Department of Neurology, Georg-August-University, Göttingen, Germany

^c DFG Research Center for Molecular Physiology of the Brain (CMPB), Göttingen, Germany

Experimental Neurology 2006 Nov;202(1):217-24

Effects of commissural de- and remyelination on motor skill behaviour in the cuprizone mouse model of multiple sclerosis

David Liebetanz^a, Doron Merkler^{b,c}

^a Department of Clinical Neurophysiology Georg-August-University, Göttingen, Germany

^b Department of Neuropathology, Georg-August-University, Robert-Koch Strasse 40, 37099 Göttingen, Germany

^c Institut für Multiple-Sklerose-Forschung, Bereich Humanmedizin der Georg-August Universität Göttingen und Gemeinnützige Hertie-Stiftung, Göttingen, Germany